



## The Pterosaur Database

### Coloborhynchus piscator

Known from several fragmentary and uncrushed fossil bones, also partial skeletons. This pterosaur from the Santana Formation of Brazil has an estimated wingspan of over 4m and a skull length of about 50cm.

**Holotype:** NSM.PV.19892  
re-classified as *Coloborhynchus piscator* in 2003 by Veldmeijer



NSM-PV.19892

Kellner, A. W. A. and Tomida., Y. **2000**, Description of a new species of Anhangueridae (Pterodactyloidea) with comments on the pterosaur fauna from the Santana Formation (Aptian -Albian), Northeastern Brazil. National Science Museum Monographs No. 17, National Science Museum, Tokyo. p. 1-135.

Veldmeijer, A. J., **2003**. Description of *Coloborhynchus spilberg* sp. nov. (Pterodactyloidea) from the Albian (Lower Cretaceous) of Brazil. *Scripta Geologica* 125: 35-139.

Veldmeijer, A. J., **2003**. Preliminary description of a skull and wing of a Brazilian Cretaceous (Santana Formation; Aptian-Albian) pterosaur (Pterodactyloidea) in the collection of the AMNH. [www.palarch.nl](http://www.palarch.nl), *vertebrate palaeontology* 0, 0: 1-13